

Standard queueing models

$Y_t \sim$ Poisson random variable with parameter λt :

$$p_{Y_t}(n) = \frac{e^{-\lambda t} (\lambda t)^n}{n!} \quad \text{for } n = 0, 1, 2, \dots \quad E[Y_t] = \lambda t$$

M/M/ ∞ :

- Steady-state probabilities: $\pi_j = \Pr\{L = j\}$ where L is a Poisson random variable with parameter λ/μ

M/M/ s :

- Steady-state probabilities:

$$\rho = \frac{\lambda}{s\mu} \quad \pi_0 = \left[\left(\sum_{j=0}^s \frac{(s\rho)^j}{j!} \right) + \frac{s^s \rho^{s+1}}{s!(1-\rho)} \right]^{-1} \quad \pi_j = \begin{cases} \frac{(\lambda/\mu)^j}{j!} \pi_0 & \text{for } j = 1, 2, \dots, s \\ \frac{(\lambda/\mu)^j}{s!s^{j-s}} \pi_0 & \text{for } j = s+1, s+2, \dots \end{cases}$$

- Expected number of customers in queue: $\ell_q = \frac{\pi_s \rho}{(1-\rho)^2}$
- Expected number of customers in the system: $\ell = \ell_q + \frac{\lambda}{\mu}$

G/G/ s :

- Whitt's approximation:

G = generic interarrival time random variable with rate $\lambda = \frac{1}{E[G]}$

X = generic service time random variable with rate $\mu = \frac{1}{E[X]}$

$$\varepsilon_a = \frac{\text{Var}[G]}{E[G]^2} \quad \varepsilon_s = \frac{\text{Var}[X]}{E[X]^2} \quad \hat{w}_q \approx \frac{\varepsilon_a + \varepsilon_s}{2} w_q$$