


**Solutions to Problem 1.** Let  $\mathcal{R} = \{2, 3, 4\}$ . Note that

$$\mathbf{P}_{\mathcal{R}\mathcal{R}} = \begin{bmatrix} 0 & 1 & 0 \\ 0.5 & 0.4 & 0.1 \\ 1 & 0 & 0 \end{bmatrix}$$

is the transition probability matrix of a self-contained Markov chain, and no proper subset of  $\mathcal{R}$  also forms a Markov chain. Therefore, states 2, 3, and 4 are recurrent, and state 1 is transient.

**Solutions to Problem 2.**


- a. From the transition probability diagram (below), we see that all states communicate with each other, because we can find a sequence of positive probability transitions that starts at state 1, goes through all the other states, and then returns to state 1 (e.g.,  $1 - 2 - 3 - 4 - 2 - 5 - 1$ ). Therefore the entire state space  $\mathcal{M} = \{1, 2, 3, 4, 5\}$  is a recurrent class, and so all states are recurrent.




- b. Let  $\mathcal{R} = \{2, 5\}$ . From the transition probability matrix, we see that  $\mathcal{R}$  is a recurrent class because

$$\mathbf{P}_{\mathcal{R}\mathcal{R}} = \begin{bmatrix} 0.8 & 0.2 \\ 0.5 & 0.5 \end{bmatrix},$$

indicating that  $\mathcal{R}$  is a self-contained Markov chain and no proper subset of  $\mathcal{R}$  is a Markov chain. Therefore, states 2 and 5 are recurrent. From the transition probability diagram (below), we can also see that states 1, 3, and 4 are transient, because the process will eventually leave each of these states and never return.



- c. From the transition probability diagram (below), we see that all states communicate with each other, because we can find a sequence of positive probability transitions that starts at state 1, goes through all the other states, and then returns to state 1 (e.g.,  $1 - 3 - 4 - 2 - 5 - 1$ ). Therefore the entire state space  $\mathcal{M} = \{1, 2, 3, 4, 5\}$  is a recurrent class, and so all states are recurrent.



**Solutions to Problem 3.** Note that states 2, 5, and 6 are absorbing states, because  $p_{ii} = 1$  for  $i = 2, 5, 6$ . In addition, note that the other states, 1, 3, and 4, are transient, because each of these states transitions to one of the absorbing states with positive probability. So,  $\mathcal{T} = \{1, 3, 4\}$ , and

$$\mathbf{N} = (\mathbf{I} - \mathbf{P}_{\mathcal{T}\mathcal{T}})^{-1} = \left( \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 0 & 0 & 0.4 \\ 0.2 & 0.4 & 0.3 \\ 0 & 0 & 0 \end{bmatrix} \right)^{-1} \approx \begin{bmatrix} 1 & 0 & 0.4 \\ 0.333 & 1.667 & 0.633 \\ 0 & 0 & 1 \end{bmatrix}$$

The absorption probabilities when  $\mathcal{R} = \{2\}$  are:

$$\alpha_{\mathcal{T}\mathcal{R}} = \mathbf{N}\mathbf{P}_{\mathcal{T}\mathcal{R}} \approx \begin{bmatrix} 1 & 0 & 0.4 \\ 0.333 & 1.667 & 0.633 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0.5 \\ 0 \\ 0.9 \end{bmatrix} \approx \begin{bmatrix} 0.86 \\ 0.737 \\ 0.9 \end{bmatrix}$$

So  $\alpha_{32} \approx 0.737$ . Similarly, the absorption probabilities when  $\mathcal{R} = \{5\}$  are:

$$\alpha_{\mathcal{T}\mathcal{R}} = \mathbf{N}\mathbf{P}_{\mathcal{T}\mathcal{R}} \approx \begin{bmatrix} 1 & 0 & 0.4 \\ 0.333 & 1.667 & 0.633 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0.1 \\ 0 \\ 0.1 \end{bmatrix} \approx \begin{bmatrix} 0.14 \\ 0.097 \\ 0.1 \end{bmatrix}$$

So  $\alpha_{35} \approx 0.097$ . Finally, the absorption probabilities when  $\mathcal{R} = \{6\}$  are:

$$\alpha_{\mathcal{T}\mathcal{R}} = \mathbf{N}\mathbf{P}_{\mathcal{T}\mathcal{R}} \approx \begin{bmatrix} 1 & 0 & 0.4 \\ 0.333 & 1.667 & 0.633 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 0.1 \\ 0 \end{bmatrix} \approx \begin{bmatrix} 0 \\ 0.167 \\ 0 \end{bmatrix}$$

So  $\alpha_{36} \approx 0.167$ .

The expected times to absorption from  $\mathcal{T}$  are:

$$\mu_{\mathcal{T}} = \mathbf{N}\mathbf{1} \approx \begin{bmatrix} 1 & 0 & 0.4 \\ 0.333 & 1.667 & 0.633 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1.4 \\ 2.633 \\ 1 \end{bmatrix}$$

Therefore,  $\mu_3 = 2.633$ .

**Solutions to Problem 4.** Looking at the transition probability matrix, we see that the Markov chain is irreducible. Let  $\mathcal{R} = \{1, 2, 3, 4\}$ . We want  $\pi_4$ :

$$\begin{aligned} \pi_{\mathcal{R}}^{\top} \mathbf{P}_{\mathcal{R}\mathcal{R}} = \pi^{\top} \\ \pi_{\mathcal{R}}^{\top} \mathbf{1} = 1 \end{aligned} \Leftrightarrow \begin{aligned} 0.70\pi_1 + 0.14\pi_2 + 0.14\pi_3 + 0.05\pi_4 &= \pi_1 \\ 0.14\pi_1 + 0.70\pi_2 + 0.14\pi_3 + 0.05\pi_4 &= \pi_2 \\ 0.14\pi_1 + 0.14\pi_2 + 0.70\pi_3 + 0.05\pi_4 &= \pi_3 \\ 0.02\pi_1 + 0.02\pi_2 + 0.02\pi_3 + 0.85\pi_4 &= \pi_4 \\ \pi_1 + \pi_2 + \pi_3 + \pi_4 &= 1 \end{aligned} \Leftrightarrow \begin{aligned} -0.30\pi_1 + 0.14\pi_2 + 0.14\pi_3 + 0.05\pi_4 &= 0 \\ 0.14\pi_1 - 0.30\pi_2 + 0.14\pi_3 + 0.05\pi_4 &= 0 \\ 0.14\pi_1 + 0.14\pi_2 - 0.30\pi_3 + 0.05\pi_4 &= 0 \\ \pi_1 + \pi_2 + \pi_3 + \pi_4 &= 1 \end{aligned}$$

Note that we removed the last equation from  $\pi_{\mathcal{R}}^{\top} \mathbf{P}_{\mathcal{R}\mathcal{R}} = \pi^{\top}$ , because any one of them is redundant. Solving this system of equations, we get:

$$\pi_1 \approx 0.2941 \quad \pi_2 \approx 0.2941 \quad \pi_3 \approx 0.2941 \quad \pi_4 \approx 0.1176$$

Therefore, the long-term market share for Poisson Puffs is 11.76%.

**Solutions to Problem 5.** Looking at the transition probability matrix, we see that the Markov chain is irreducible. Let  $\mathcal{R} = \{1, 2, 3, 4\}$ . We want  $\pi_4$ :

$$\begin{aligned} \pi_{\mathcal{R}}^{\top} \mathbf{P}_{\mathcal{R}\mathcal{R}} = \pi^{\top} \\ \pi_{\mathcal{R}}^{\top} \mathbf{1} = 1 \end{aligned} \Leftrightarrow \begin{aligned} 0\pi_1 + \frac{1}{3}\pi_2 + \frac{1}{3}\pi_3 + \frac{1}{3}\pi_4 &= \pi_1 \\ \frac{1}{2}\pi_1 + 0\pi_2 + \frac{1}{3}\pi_3 + \frac{1}{3}\pi_4 &= \pi_2 \\ \frac{1}{2}\pi_1 + \frac{1}{3}\pi_2 + 0\pi_3 + \frac{1}{3}\pi_4 &= \pi_3 \\ 0\pi_1 + \frac{1}{3}\pi_2 + \frac{1}{3}\pi_3 + 0\pi_4 &= \pi_4 \\ \pi_1 + \pi_2 + \pi_3 + \pi_4 &= 1 \end{aligned} \Leftrightarrow \begin{aligned} -\pi_1 + \frac{1}{3}\pi_2 + \frac{1}{3}\pi_3 + \frac{1}{3}\pi_4 &= 0 \\ \frac{1}{2}\pi_1 - \pi_2 + \frac{1}{3}\pi_3 + \frac{1}{3}\pi_4 &= 0 \\ \frac{1}{2}\pi_1 + \frac{1}{3}\pi_2 - \pi_3 + \frac{1}{3}\pi_4 &= 0 \\ \pi_1 + \pi_2 + \pi_3 + \pi_4 &= 1 \end{aligned}$$

Note that we removed the last equation from  $\pi_{\mathcal{R}}^{\top} \mathbf{P}_{\mathcal{R}\mathcal{R}} = \pi^{\top}$ , because any one of them is redundant. Solving this system of equations, we get:

$$\pi_1 = \frac{1}{4} \quad \pi_2 = \frac{9}{32} \quad \pi_3 = \frac{9}{32} \quad \pi_4 = \frac{3}{16}$$

Therefore, the AGV spends 3/16 of the time at the output buffer in the long run.

**Solutions to Problem 6.**

- Looking at the transition probability diagram, we can see that  $\{1, 2\}$  and  $\{3, 5\}$  form self-contained Markov chains, and no proper subsets of  $\{1, 2\}$  or  $\{3, 5\}$  form a self-contained Markov chain.
- Recurrent states: 1, 2, 3, 5 (these are states that are part of a recurrent class, by part a)  
Transient states: 4, 6 (these are states not part of a recurrent class)


- Let  $\mathcal{R} = \{1, 2\}$ . We want  $\pi_1$ . From the transition probability diagram,  $\mathbf{P}_{\mathcal{R}\mathcal{R}} = \begin{bmatrix} 0.2 & 0.8 \\ 0.7 & 0.3 \end{bmatrix}$ . Therefore,

$$\begin{aligned} \pi_{\mathcal{R}}^{\top} \mathbf{P}_{\mathcal{R}\mathcal{R}} = \pi^{\top} \\ \pi_{\mathcal{R}}^{\top} \mathbf{1} = 1 \end{aligned} \Leftrightarrow \begin{aligned} 0.2\pi_1 + 0.7\pi_2 &= \pi_1 \\ 0.8\pi_1 + 0.3\pi_2 &= \pi_2 \\ \pi_1 + \pi_2 &= 1 \end{aligned} \Rightarrow \pi_1 = \frac{7}{15}, \pi_2 = \frac{8}{15}$$

So, the long-run fraction of time the UAV spends in region 1 is 7/15.

- d. This is a little tricky: the definition of an absorbing probability requires an absorbing state, that is, a recurrent class with only one state.

Let's replace states 3 and 5 with a "super state" called 7. We end up with the following transition probability diagram:



Now, let  $\mathcal{T} = \{4, 6\}$  and  $\mathcal{R} = \{7\}$ . We want  $\alpha_{47}$ :

$$\alpha_{\mathcal{T}\mathcal{R}} = (\mathbf{I} - \mathbf{P}_{\mathcal{T}\mathcal{T}})^{-1} \mathbf{P}_{\mathcal{T}\mathcal{R}} = \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 0.1 & 0.2 \\ 0.1 & 0.1 \end{pmatrix} \right)^{-1} \begin{bmatrix} 0.5 \\ 0.7 \end{bmatrix} \approx \begin{bmatrix} 0.747 \\ 0.861 \end{bmatrix}$$

Therefore,  $\alpha_{47} \approx 0.747$ .