## Quiz 1 - 8/23/2023

**Instructions.** You have 15 minutes to complete this quiz. You may use your plebe-issue TI-36X Pro calculator. You may <u>not</u> use any other materials.

<u>Show all your work.</u> To receive full credit, your solutions must be completely correct, sufficiently justified, and easy to follow.

Problem la	Weight	Score
1b	1	
2a	1	
2b	1	
Total		/ 40

**Problem 1.** Simplex Pizza sells New York style and Sicilian style pizza by the slice. Let N represent the number of New York style slices in one order, and let T represent the  $\underline{\text{total}}$  number of slices in one order. The joint probabilities of N and T are as follows:

			T	
		1	2	3
N	0	0.10	0.05	0.01
	1	0.25	0.10	0.02
	2	0	0.35	0.04
	3	0	0	0.08

a. What is the probability that an order contains a total of 2 slices?

b. What is the probability that an order contains 2 New York style slices, given that the order contains a total of 2 slices?

Problem 2. The Orange Company was having problems with its automated manufacturing cells yesterday: sometim	es
a tablet came out of a cell defective. 65% of the tablets were produced in cell 1, 25% in cell 2, and 10% in cell 3. 1% of the	hε
tablets produced in cell 1 came out defective, 1% in cell 2, and 5% in cell 3.	

Suppose you select 1 tablet made yesterday at random. Let C be a random variable that represents the cell it was produced in (i.e., C = 1, 2 or 3). In addition, let D represent a random variable indicating whether the tablet came out defective (i.e., D = 1 if defective, 0 otherwise).

a. What is the probability that the randomly selected tablet came out defective, i.e.  $Pr\{D = 1\}$ ?

b. Are *C* and *D* independent? Give a numerical argument for why or why not.