

Lesson 11. Derivatives and Integrals of Vector Functions

0 Warm up

Example 1. Find these derivatives and integrals.

(a) $\frac{d}{dt}(1+t^3) =$

(d) $\int 2t \, dt =$

(b) $\frac{d}{dt}(\cos 2t) =$

(e) $\int 2 \cos t \, dt =$

(c) $\frac{d}{dt}(te^{-t}) =$

(f) $\int_0^{2\pi} 3 \, dt =$

1 Today...



- Derivatives of vector functions
- Integrals of vector functions
- Arc length

2 Derivatives

- The **derivative** of \vec{r} is

$$\frac{d\vec{r}}{dt} = \vec{r}'(t) = \lim_{h \rightarrow 0} \frac{\vec{r}(t+h) - \vec{r}(t)}{h}$$

- Note: the derivative of a vector function is also a vector



- Let C be the curve defined by \vec{r}
- Let $\vec{r}(t)$ be the position vector of P
- The derivative $\vec{r}'(t)$ is the direction vector of the line tangent to C at P
 \Rightarrow Sometimes we refer to $\vec{r}'(t)$ as the **tangent vector**


- The **unit tangent vector** is

- If $\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$, where f , g , and h are differentiable functions, then

Example 2.

- (a) Find the derivative of $\vec{r}(t) = \langle \cos 2t, 1 + t^3, te^{-t} \rangle$.
- (b) Find the unit tangent vector at the point where $t = 0$.

Example 3. Find parametric equations for the tangent line to the curve given by $\vec{r}(t) = \langle \cos t, \sin t, t \rangle$ at point $(0, 1, \pi/2)$.



- Differentiation rules

$$1. \frac{d}{dt}(\vec{u}(t) + \vec{v}(t)) = \vec{u}'(t) + \vec{v}'(t)$$

$$4. \frac{d}{dt}(\vec{u}(t) \cdot \vec{v}(t)) = \vec{u}'(t) \cdot \vec{v}(t) + \vec{u}(t) \cdot \vec{v}'(t)$$

$$2. \frac{d}{dt}(c\vec{u}(t)) = c\vec{u}'(t)$$

$$5. \frac{d}{dt}(\vec{u}(t) \times \vec{v}(t)) = \vec{u}'(t) \times \vec{v}(t) + \vec{u}(t) \times \vec{v}'(t)$$

$$3. \frac{d}{dt}(f(t)\vec{u}(t)) = f'(t)\vec{u}(t) + f(t)\vec{u}'(t)$$

$$6. \frac{d}{dt}(\vec{u}(f(t))) = f'(t)\vec{u}'(f(t))$$

3 Integration

- Let f , g , and h be continuous functions
- The **indefinite integral** of a vector function $\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$ is

- The **definite integral** of a vector function $\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$ from a to b is

- Note: the integral of a vector function is also a vector

Example 4. Let $\vec{r}(t) = \langle 2 \sin t, 2 \cos t, 2t \rangle$. Find $\int_0^{\pi/2} \vec{r}(t) dt$.

4 Arc length

- Let C be a curve with vector equation $\vec{r}(t) = \langle f(t), g(t), h(t) \rangle$, $a \leq t \leq b$
- What is the length of C ?
- The **arc length** of C is

- Similar for curves in 2D

Example 5. Let C be the curve defined by $\vec{r}(t) = \langle \cos t, \sin t, t \rangle$, $0 \leq t \leq 2\pi$. Find the length of C .

