

Lesson 2. Vectors

1 Today...

- Vectors in \mathbb{R}^2 and \mathbb{R}^3
- Standard basis vectors and unit vectors
- Problems with forces

2 What is a vector?


- A **vector** is an object that has both

and

- Notation: \mathbf{a} or \vec{a}
- A vector \vec{a} can be represented by an ordered list of numbers:

$$\vec{a} = \langle a_1, a_2 \rangle \quad (\text{in } \mathbb{R}^2) \qquad \vec{a} = \langle a_1, a_2, a_3 \rangle \quad (\text{in } \mathbb{R}^3)$$

- These numbers (e.g. a_1, a_2, a_3) are known as **components** of \vec{a}
- For now, let's stick to \mathbb{R}^2
 - Much of what we'll see generalizes to \mathbb{R}^3
- Graphically:




- The **magnitude** or **length** of vector $\vec{a} = \langle a_1, a_2 \rangle$ is


- Two vectors are **equivalent** if they have the same magnitude and direction – position does not matter
- A special vector – the **zero vector** $\vec{0} = \langle 0, 0 \rangle$
 - $\vec{0}$ is the only vector with no specific direction

Example 1. Consider the vectors below.

- Give \vec{a} , \vec{b} , \vec{c} as an ordered list of numbers.
- Find $|\vec{c}|$.
- Are any of these vectors equivalent? Which ones?



Example 2. Consider \vec{d} below. We are given that $|\vec{d}| = 5$. Give \vec{d} as an ordered list of numbers.



3 Scalar multiplication

- Let c be a scalar, \vec{a} be a vector $\Rightarrow c\vec{a}$ is another vector


- $c\langle a_1, a_2 \rangle =$

Example 3. Consider $\vec{a} = \langle 2, 1 \rangle$ from the previous example.

a. $3\vec{a} =$

b. $-2\vec{a} =$

- c. Draw \vec{a} , $3\vec{a}$ and $-2\vec{a}$ below.



- If $c > 0$, then $c\vec{a}$ is a vector in the same direction as \vec{a} and $|c|$ times the length of \vec{a}
- If $c < 0$, then $c\vec{a}$ is a vector in the opposite direction as \vec{a} and $|c|$ times the length of \vec{a}
- If $c = 0$, then $c\vec{a} = \vec{0}$

4 Adding and subtracting vectors



- Let \vec{a}, \vec{b} be vectors $\Rightarrow \vec{a} + \vec{b}$ is another vector

- $\langle a_1, a_2 \rangle + \langle b_1, b_2 \rangle =$


Example 4. Consider $\vec{a} = \langle 2, 1 \rangle$ and $\vec{b} = \langle 1, 2 \rangle$ from the previous example.

a. $\vec{a} + \vec{b} =$


b. Draw $\vec{a} + \vec{b}$ below:



- **Triangle law** for adding vectors:



- **Parallelogram law** for adding vectors:



- $\langle a_1, a_2 \rangle - \langle b_1, b_2 \rangle =$

5 Generalizations

- All of the above generalizes naturally to \mathbb{R}^3 :

$$\begin{aligned} |\langle a_1, a_2, a_3 \rangle| &= \sqrt{a_1^2 + a_2^2 + a_3^2} & \langle a_1, a_2, a_3 \rangle + \langle b_1, b_2, b_3 \rangle &= \langle a_1 + b_1, a_2 + b_2, a_3 + b_3 \rangle \\ c \langle a_1, a_2, a_3 \rangle &= \langle ca_1, ca_2, ca_3 \rangle & \langle a_1, a_2, a_3 \rangle - \langle b_1, b_2, b_3 \rangle &= \langle a_1 - b_1, a_2 - b_2, a_3 - b_3 \rangle \end{aligned}$$

- Algebraically, vectors behave a lot like scalars, e.g.

$$\vec{a} + \vec{b} = \vec{b} + \vec{a} \quad c(\vec{a} + \vec{b}) = c\vec{a} + c\vec{b} \quad (c + d)\vec{a} = c\vec{a} + d\vec{a}$$

- See p. 802 of Stewart for a fuller list

6 Standard basis vectors and unit vectors

- **Standard basis vectors** in \mathbb{R}^3 :

$$\vec{i} = \boxed{} \quad \vec{j} = \boxed{} \quad \vec{k} = \boxed{}$$

- We can write any vector as the sum of scalar multiples of standard basis vectors:

- A **unit vector** is a vector with length 1

◦ For example, $\vec{i}, \vec{j}, \vec{k}$ are all unit vectors

- The unit vector that has the same direction as \vec{a} (assuming $\vec{a} \neq \vec{0}$) is

Example 5. Let $\vec{a} = 4\vec{i} - \vec{j} + 2\vec{k}$ and $\vec{b} = \vec{i} + 2\vec{k}$.


- Write $\vec{a} - 2\vec{b}$ in terms of $\vec{i}, \vec{j}, \vec{k}$.
- Find a unit vector in the direction of $\vec{a} - 2\vec{b}$.

- Note: all of this applies to vectors in \mathbb{R}^2 in a similar way

7 Problems with forces

- Some physics:
 - **Force** has magnitude and direction, and so it can be represented by a vector
 - Force is measured in pounds (lbs) or newtons (N)
 - If several forces are acting on an object, the **resultant force** experienced by the object is the sum of these forces

Example 6. A weight \vec{w} counterbalances the tensions (forces) in two wires as shown below:



The tensions \vec{T}_1 and \vec{T}_2 both have a magnitude of 20lb. Find the magnitude of the weight \vec{w} .

- Note: if an object has a mass of m kg, then it has a weight of mg N, where $g = 9.8$