

Lesson 20. Tangent Planes and Linear Approximations

0 Warm up


Example 1. Find an equation of the plane that passes through $(-1, 1, 5)$ and is perpendicular to the vector $\langle 2, 4, -3 \rangle$.

1 Tangent planes

- Let S be a surface with equation $z = f(x, y)$
- Let $P(x_0, y_0, z_0)$ be a point on S

◦ So $z_0 =$

- Let T_1 and T_2 be the tangent lines at P in the x - and y -directions, respectively
- The **tangent plane** to the surface S at point P is the plane that contains both tangent lines T_1 and T_2



- The tangent plane must have an equation of the form $a(x - x_0) + b(y - y_0) + c(z - z_0) = 0$ or equivalently,

- If this is the equation of the tangent plane, its intersection with the plane $y = y_0$ must be the tangent line T_1

• Setting $y = y_0$, we obtain

- Looking at this equation a must be

• Similarly, b must be

⇒ An equation of the tangent plane to the surface $z = f(x, y)$ at point $P(x_0, y_0, z_0)$ is




Example 2. Find the tangent plane to the surface $z = x^2 + xy + 3y^2$ at the point $(1, 1, 5)$.

2 Linear approximations


- What do the level curves of a plane look like?

- As we zoom in on the level curves of an arbitrary surface, they start to look more and more like equally spaced parallel lines


- For example: $f(x, y) = 2x^2 + y^2$




⇒ We can use tangent planes to approximate function values



(a)



(b)



(c)

- The **linear approximation** of f at (a, b) is

- Compare to equation for tangent plane above: use $x_0 = a$, $y_0 = b$, $z_0 = f(a, b)$

Example 3. Find the linear approximation of $f(x, y) = xe^{xy}$ at $(1, 0)$. Use it to approximate $f(1.1, -0.2)$.

Example 4. Here is the wind-chill index function $W(T, v)$ we have seen in previous lessons:

		Wind speed (km/h)											
		5	10	15	20	25	30	40	50	60	70	80	
Actual temperature (°C)	$T \backslash v$	5	4	3	2	1	1	0	-1	-1	-2	-2	-3
	0	-2	-3	-4	-5	-6	-6	-7	-8	-9	-9	-10	-10
	-5	-7	-9	-11	-12	-12	-13	-14	-15	-16	-16	-17	-17
	-10	-13	-15	-17	-18	-19	-20	-21	-22	-23	-23	-24	-24
	-15	-19	-21	-23	-24	-25	-26	-27	-29	-30	-30	-31	-31
	-20	-24	-27	-29	-30	-32	-33	-34	-35	-36	-36	-37	-38
	-25	-30	-33	-35	-37	-38	-39	-41	-42	-43	-43	-44	-45
	-30	-36	-39	-41	-43	-44	-46	-48	-49	-50	-50	-51	-52
	-35	-41	-45	-48	-49	-51	-52	-54	-56	-57	-57	-58	-60
	-40	-47	-51	-54	-56	-57	-59	-61	-63	-64	-64	-65	-67

Once upon a time, we estimated $W_T(-15, 40) \approx 1.3$. In a similar fashion, we can estimate $W_v(-15, 40) \approx -0.15$.

Find the linear approximation of $W(T, v)$ at $(-15, 40)$. Use it to approximate $W(-12, 45)$.

- Why bother with linear approximations?
 - Desert island
 - More importantly: **linear functions** (functions of the form $f(x, y) = ax + by$) are much easier to deal with than other types of functions
 - ⇒ Linear approximations form the basis of many algorithms for complex problems
- Disclaimer: equations for tangent planes and linear approximations above do not necessarily apply when the partial derivatives of f are not continuous