

Lesson 30. Double Integrals over Rectangles

0 Warm up

Example 1. Find the value of


a. $\sum_{i=2}^4 i =$

b. $\sum_{i=1}^3 \sum_{j=1}^2 ij =$


1 Review: area and integrals

- The definite integral of a single-variable function:

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$$



2 Volume and double integrals




- Let R be a rectangle in the xy -plane:

$$R = [a, b] \times [c, d] = \{(x, y) : a \leq x \leq b, c \leq y \leq d\}$$

- Let $f(x, y)$ be a function of two variables
- What is the volume of the solid above R and below the graph of f ?

- Idea:
 - Divide R into subrectangles of equal area ΔA
 - ◊ Grid with m columns (x -direction) and n rows (y -direction)
 - For each subrectangle R_{ij} :
 - ◊ Choose a **sample point** (x_{ij}^*, y_{ij}^*)
 - ◊ Compute the volume of the (thin) rectangular box with base R_{ij} and height $f(x_{ij}^*, y_{ij}^*)$.
 - Add the volumes of all these rectangular boxes



- Estimated volume:

- This is called a **double Riemann sum**

- The **double integral** of f over the rectangle R is


- How do we choose sample points in each subrectangle?

- Upper right corner
- Lower left corner
- **Midpoint rule:** center of subrectangle

- If $f(x, y) \geq 0$, then the volume V of the solid that lies above the rectangle R and below the surface $z = f(x, y)$ is

Example 2. Estimate the volume of the solid that lies above the square $R = [0, 2] \times [0, 2]$ and below $f(x, y) = 16 - x^2 - 2y^2$. Use a Riemann sum with $m = 2$ and $n = 2$. Use the upper right corners as sample points.

Example 3. Below is a contour map for a function f on the square $R = [0, 3] \times [0, 3]$. Use a Riemann sum with $m = 3$ and $n = 3$ to estimate the value of $\iint_R f(x, y) dA$. Use the midpoint rule to take sample points.



3 Average value

- The **average value** of a function of two variables defined on a rectangle R is

Example 4. Estimate the average value of the function f in Example 3 on R .

4 Iterated integrals

- **Partial integration** with respect to x : $\int_a^b f(x, y) dx$
 - Regard y as a constant (i.e., fixed, coefficient, etc.)
 - Integrate $f(x, y)$ with respect to x from $x = a$ to $x = b$
 - Results in an expression in terms of y
- Partial integration with respect to y defined in a similar way
- **Iterated integrals**: work from the inside out
 - $\int_c^d \int_a^b f(x, y) dx dy = \int_c^d \left[\int_a^b f(x, y) dx \right] dy$
 - ◊ Integrate first with respect to x from $x = a$ to $x = b$ (keeping y constant)
 - ◊ Integrate resulting expression in y with respect to y from $y = c$ to $y = d$
 - $\int_a^b \int_c^d f(x, y) dy dx = \int_a^b \left[\int_c^d f(x, y) dy \right] dx$
 - ◊ Integrate first with respect to from
 - ◊ Integrate resulting expression in with respect to from

Example 5. Evaluate $\int_0^3 \int_1^2 x^2 y dy dx$.

Example 6. Evaluate $\int_1^2 \int_0^3 x^2 y dx dy$.

- **Fubini's theorem for rectangles.** If $R = [a, b] \times [c, d]$, then:

- (f needs to satisfy some conditions, e.g. f is continuous on R)
- Double integrals over rectangles can be evaluated using iterated integrals
- Order of integration does not matter!

Example 7. Evaluate $\iint_R (x - 3y^2) dA$, where $R = [0, 2] \times [1, 2]$.

Example 8. Find the volume of the solid that is bounded by the surface $x^2 + 2y^2 + z = 16$, the planes $x = 2$ and $y = 2$, and the three coordinate planes.

Example 9. Evaluate $\int_0^1 \int_0^1 ye^{xy} dy dx$.