


## Lesson 31. Double Integrals Over General Regions

### 1 Last time: rectangles




- Rectangle  $R = [a, b] \times [c, d]$   
 $= \{(x, y) \mid a \leq x \leq b, c \leq y \leq d\}$

- $$\iint_R f(x, y) dA = \int_a^b \int_c^d f(x, y) dy dx$$

$$= \int_c^d \int_a^b f(x, y) dx dy$$



### 2 General regions



- How about general regions  $D$ ?
- Intuition: if  $f(x, y) \geq 0$ , double integral still represents volume of solid between  $D$  and graph of  $f$
- We focus on two types of regions

- **Type I regions:** lies between the graphs of two continuous functions of  $x$ , that is

- Examples:





- If  $D$  is a type I region (and  $f$  is continuous on  $D$ ), then

- In the inner integral,  $x$  is regarded a constant in  $f(x, y)$  and the limits of integration

- **Type II regions:** lies between the graphs of two continuous functions of  $y$ , that is

- Examples:



- If  $D$  is a type II region (and  $f$  is continuous on  $D$ ), then


- In the inner integral,  $y$  is regarded a constant in  $f(x, y)$  and the limits of integration

**Example 1.** Find  $\iint_D (x - y) dy dx$ , where  $D = \{(x, y) \mid 0 \leq x \leq 1, 2x \leq y \leq 2\}$ .

**Example 2.** Find  $\iint_D (x + 2) dA$ , where  $D$  is the region bounded by the parabolas  $y = 2x^2$  and  $y = 1 + x^2$ .




**Example 3.** Evaluate  $\iint_D (2 - 2x - y) dA$ , where  $D$  is the triangular region with vertices  $(0, 0)$ ,  $(2, 0)$ ,  $(1, 1)$ .




- If  $D = D_1 \cup D_2$ , where  $D_1$  and  $D_2$  don't overlap except perhaps on their boundaries, then

$$\iint_D f(x, y) dA = \iint_{D_1} f(x, y) dA + \iint_{D_2} f(x, y) dA$$




(a)  $D$  is neither type I nor type II.




(b)  $D = D_1 \cup D_2$ ,  $D_1$  is type I,  $D_2$  is type II.

**Example 4.** Write  $\iint_D (2 - 2x - y) dA$  as the sum of 2 type I region iterated integrals, where  $D$  is the triangular region with vertices  $(0, 0)$ ,  $(2, 0)$ , and  $(1, 1)$ .

**Example 5.** Sketch the region of integration of  $\int_0^2 \int_{x^2}^{2x} (x^2 + y^2) dy dx$ . Change the order of integration. Evaluate the integral using the order of integration of your choice.



**Example 6.** Consider the double integral  $\iint_D f(x, y) dA$  where  $D$  is enclosed by  $x = 0$ ,  $x = \sqrt{1 - y^2}$ . Set up this double integral as an iterated integral using both orders of integration.



**Example 7.** Consider the double integral  $\int_0^4 \int_{\sqrt{x}}^2 f(x, y) dy dx$ . Sketch the region of integration and change the order of integration.



**Example 8.** Let  $D$  be some region in the  $xy$ -plane. What does  $\iint_D 1 dA$  represent? Explain.

