

## Lesson 15. The Chain Rule

### 1 This lesson

- Two special cases of the chain rule
- Tree diagrams and the general version of the chain rule

### 2 Case 1

- Let  $z = f(x, y)$  be a function of 2 variables
- Let  $x$  and  $y$  be functions of 1 variable:  $x = g(t)$  and  $y = h(t)$   
 $\Rightarrow z$  is indirectly a function of  $t$ :  $z = f(g(t), h(t))$
- Can we find the derivative of  $z$  with respect to  $t$ ?

- Chain rule (Case 1):

**Example 1.** Let  $z = xy - x^2y$ ,  $x = t^2 + 1$ , and  $y = t - 1$ . Find  $dz/dt$ .

**Example 2.** Let  $z = x^2y + 3xy^4$ ,  $x = \sin 2t$ , and  $y = \cos t$ . Find  $dz/dt$  when  $t = 0$ .

**Example 3.** A person's body-mass index (BMI) is given by  $B(h, w) = 700w/h^2$ , where  $h$  is the person's height in inches and  $w$  is the person's weight in pounds.

Suppose MIDN Slim is currently 70 inches tall and 140 pounds. MIDN Slim is growing at a rate of 0.1 inches per year, and is gaining weight at a rate of 2 pounds per year. Find the rate at which MIDN Slim's BMI is changing per year.

### 3 Case 2

- Let  $z = f(x, y)$  be a function of 2 variables
- Let  $x$  and  $y$  be functions of 2 variables:  $x = g(s, t)$  and  $y = h(s, t)$   
 $\Rightarrow z$  is indirectly a function of  $s$  and  $t$ :  $z = f(g(s, t), h(s, t))$
- We can find the derivative of  $z$  with respect to  $s$  and  $t$
- **Chain rule (Case 2):**


**Example 4.** Let  $z = \sin x \cos y$ ,  $x = st^2$ ,  $y = s^2t$ . Find  $\partial z/\partial s$  and  $\partial z/\partial t$ .

**Example 5.** Suppose  $f$  is a differentiable function of  $x$  and  $y$ , and  $g(u, v) = f(e^u + \sin v, e^u + \cos v)$ . Use the following table to compute  $g_u(0, 0)$  and  $g_v(0, 0)$ .

	$f$	$g$	$f_x$	$f_y$
$(0, 0)$	1	4	8	0
$(1, 2)$	4	1	3	6

#### 4 Tree diagrams

- How can we remember the more complex chain rule (i.e. Case 2)?
- Draw a **tree diagram**:



- To get  $\partial z/\partial s$ , follow all the paths from  $z$  to  $s$ :

- This idea can be extended in general to functions of 3 or more variables

**Example 6.** Write out the chain rule for the case where  $z = f(w, x, y)$ ,  $w = g(s, t)$ ,  $x = h(s, t)$ ,  $y = \ell(s, t)$ .

**Example 7.** Write out the chain rule for the case where  $w = f(x, y, z)$ ,  $x = g(t)$ ,  $y = h(t)$ ,  $z = \ell(t)$ .

**Example 8.** Let  $w = \ln(\sqrt{x^2 + y^2 + z^2})$ ,  $x = \sin t$ ,  $y = \cos t$ ,  $z = t$ . Find  $dw/dt$ .

**Example 9.** The length  $\ell$ , width  $w$ , and height  $h$  of a box change with time. At a certain instant the dimensions are  $\ell = 1$  m,  $w = 2$  m, and  $h = 2$  m.  $\ell$  and  $w$  are increasing at a rate of 2 m/s while  $h$  is decreasing at a rate of 3 m/s. Find the rate at which the length of the diagonal is changing at that instant.