

## Lesson 24. Double Integrals in Polar Coordinates

### 1 Review


#### 1.1 Polar coordinates

- **Polar coordinate system:** specify points in the  $xy$ -plane as  $(r, \theta)$  where

◦  $r =$

◦  $\theta =$


**Example 1.** Sketch the region in the plane consisting of points whose polar coordinates satisfy:  $1 \leq r \leq 3$ ,  $\pi/6 \leq \theta \leq 5\pi/6$ .




#### 1.2 Polar curves

- The **graph of a polar equation**  $F(r, \theta) = 0$  consists of all points that can be represented by some polar coordinates  $(r, \theta)$  that satisfy the equation

**Example 2.** Sketch the curve with polar equation  $r = 2 \cos \theta$ .



### 1.3 Correspondence between polar and Cartesian coordinates



•  $x =$

•  $y =$

•

•


**Example 3.** Find a Cartesian equation for the curve  $r = 2 \cos \theta$ .

**Example 4.** Find a polar equation for the curve represented by the Cartesian equation  $4y^2 = x^2$ .


## 2 Changing to polar coordinates in a double integral

- Idea:

- Some regions are hard to express in terms of rectangular coordinates, but easily described using polar coordinates




(a)  $R = \{(r, \theta) \mid 0 \leq r \leq 1, 0 \leq \theta \leq 2\pi\}$



(b)  $R = \{(r, \theta) \mid 1 \leq r \leq 2, 0 \leq \theta \leq \pi\}$



- How do we integrate in polar coordinates? Divide regions into **polar subrectangles**



- If  $D$  is a polar region of the form

$$D = \{(r, \theta) \mid \alpha \leq \theta \leq \beta, h_1(\theta) \leq r \leq h_2(\theta)\}$$

then



- Substitute  $x = r \cos \theta$  and  $y = r \sin \theta$  into  $f(x, y)$
- Replace  $dA$  with  $r \, dr \, d\theta$
- Don't forget the additional factor  $r$ !**

**Example 5.** Evaluate  $\int_{-3}^3 \int_0^{\sqrt{9-x^2}} \sin(x^2 + y^2) dy dx$  by converting to polar coordinates.

**Example 6.** Evaluate  $\iint_D (x^2 + y^2) dA$ , where  $D$  is the region in the first quadrant bounded by the circles  $x^2 + y^2 = 1$  and  $x^2 + y^2 = 4$  and the lines  $x = 0$  and  $y = x$ .

**Example 7.** Find the volume of the solid bounded by the plane  $z = 0$  and the paraboloid  $z = 1 - x^2 - y^2$ .